



NEBRASKA CANCER SPECIALISTS

the Physicians of Oncology Hematology West

Votrient®

Generic name: Pazopanib

Drug type:

Votrient is a targeted therapy. It is classified as a kinase inhibitor.

How Votrient is given:

- Votrient is given by mouth.
- Votrient must be taken on an empty stomach, one hour before, or two hours after food.

Side effects of Votrient:

Important things to remember about the side effects of Votrient:

- Most people do not experience all of the side effects listed.
- Side effects are almost always reversible and will go away after treatment is complete.
- There are many options to help minimize or prevent side effects.

The following side effects are common for patients receiving Votrient:

- Diarrhea
- Increase in blood pressure
- Hair color changes

These are less common side effects for patients receiving Votrient:

- Nausea/vomiting
- Decreased desire to eat
- Fatigue
- Abdominal pain
- Muscle aches
- Headache
- Alterations in blood chemistry tests

When to contact your doctor or health care provider:

Contact your health care provider ***immediately***, day or night, if you should experience any of the following symptoms:

- Fever of 101° F or higher, chills (possible signs of infection)

- Difficulty breathing, sudden weight gain, swelling, vision changes
- Severe abdominal pain

The following symptoms require medical attention, but are not emergency situations. Contact your health care provider ***within 24 hours*** of noticing any of the following:

- Nausea (interferes with ability to eat and unrelieved with prescribed medications)
- Vomiting (vomiting more than 4-5 times in a 24-hour period)
- Mouth sores (painful redness, swelling or ulcers)
- Diarrhea (4-6 episodes in a 24-hour period)
- Constipation unrelieved by laxative use
- Any signs of bleeding, including black or tarry stools, or blood in your stools or urine
- Signs of infection such as redness or swelling, pain on swallowing, coughing up mucous, or painful urination
- Swelling of the feet or ankles
- Sudden weight gain
- Bone or joint pain
- Yellowing of the skin or eyes

Always inform your health care provider if you experience any unusual symptoms.